Neuroscience Letters, 128 (1991) 305–308
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Adamson, P., see Brammer, M.J., 231
Agresti, C., see Levi, G., 37
Akaoka, H., see Luppi, P.-H., 33
Aloisi, F., see Levi, G., 37
Altmann, L., Sveinsson, K. and Wiegand, H., Long-term potentiation in rat hippocampal slices is impaired following acute lead perfusion, 109
Al-Yousuf, S. and Mizuno, N., Electron microscopic identification of axons containing melanin-concentrating hormone in the lamprey, *Lampetra fluviatilis* L., 249
Anand, P., Ghatei, M.A., Christofides, N.D., Blank, M.A., McGregor, G.P., Morrison, J.F.B., Scaravilli, F. and Bloom, S.R., Differential neuropeptide expression after visceral and somatic nerve injury in the cat and rat, 57
Anderson, J.P., Esch, F.S., Keim, P.S., Sambamurti, K., Lieberburg, I. and Robakis, N.K., Exact cleavage site of Alzheimer amyloid precursor in neuronal PC-12 cells, 126
Ando, S. and Ohashi, Ya., Longitudinal study on age-related changes of working and reference memory in the rat, 17
Annies, R., see Löscher, W., 191
Anwar, M., see Park, D.H., 240
Arbuthnott, G.W., see Wright, A.K., 281

Barden, N., see L'Héreault, S., 207
Barnes, B.M., see Daan, S., 265
Bartfai, T., see Bristulf, J., 173
Bassant, M.H., see Lee, B.H., 29
Beyer, C., Pilgrim, C., Reisert, I. and Misgeld, U., Cells from embryonic rat striatum cocultured with mesencephalic glia express dopaminergic phenotypes, 1
Blanchard, J.C., see Stutzmann, J.M., 4
Blank, M.A., see Anand, P., 57
Bloch, S., see Musumeci, D., 285
Bloom, S.R., see Anand, P., 57
Bose, R., see Sutherland, G., 169
Brammer, M.J., Richmond, S., Xiang, J.Z., Adamson, P., Hajimohammadreza, I., Silva, M.A. and Campbell, I.C., Kainate and quisqualate effects on rat presynaptic cortical receptors are metabotropic and non-additive, 231
Brené, S., see O'Connor, W.T., 66
Bristulf, J., Simoncsits, A. and Bartfai, T., Characterization of a neuronal interleukin-1 receptor and the corresponding mRNA in the mouse anterior pituitary cell line AtT-20, 173
Brundin, Lou, Flock, Åke, Khanna, S.M. and Ulfendahl, M., Frequency-specific position shift in the guinea pig organ of Corti, 77
Bugiani, O., see Tagliavini, F., 117

Campbell, I.C., see Brammer, M.J., 231
Charlety, P.J., see Luppi, P.-H., 33
Chouvet, G., see Luppi, P.-H., 33
Christofides, N.D., see Anand, P., 57

Daan, S., Barnes, B.M. and Strijkstra, A.M., Warming up for sleep? — ground squirrels sleep during arousals from hibernation, 265
Darche, F., see Stutzmann, J.M., 4
Denney, R.M., see Lin, J.S., 61
Dermietzel, R., see Krause, D., 301
Dunkley, P.R., see Heath, J.W., 277
Dunnett, S.B., see Wright, A.K., 281
D'Urso, D., see Levi, G., 37

Eisen, A., see Seltzer, Z., 203
Emson, P.C., see Iwamoto, N., 81
Emson, P.C., see Kiyama, Hiroshi, 85
Eon, B., see Stutzmann, J.M., 4
Eriksson, S.V., see Lundeberg, T., 161
Esch, F.S., see Anderson, J.P., 126

Feringa, E.R., see Garrett, W.T., 137
Flock, Åke, see Brundin, Lou, 77
Fort, P., see Lin, J.S., 61
Fort, P., see Luppi, P.-H., 33
Frangione, B., see Tagliavini, F., 117
Friedel, I., see Thürauf, N., 297
Fukui, H., see Inagaki, N., 257

Galla, H.J., see Krause, D., 301
Garrett, W.T., McBride, R.L., Williams Jr., J.K. and Feringa, E.R., Fluoro-Gold's toxicity makes it inferior to True Blue for long-term studies of dorsal root ganglion neurons and motoneurons, 137
Gentle, M.J., Hunter, L.N. and Waddington, D., The onset of pain related behaviours following partial beak amputation in the chicken, 113
Ghatei, M.A., see Anand, P., 57
Ghiso, J., see Tagliavini, F., 117
Giaccone, G., see Tagliavini, F., 117
Ginzburg, R., see Seltzer, Z., 203

Hajimohammadreza, I., see Brammer, M.J., 231
Hao, J.-X., Xu, X.-J., Yu, Y.-X., Seiger, Å. and Wiesenfeld-Hallin, Z., Hypersensitivity of dorsal horn wide dynamic range neurons to cutaneous mechanical stimuli after transient spinal cord ischemia in the rat, 105
Harino, S., see Riva, C.E., 291
Hatase, O., see Yamaguchi, F., 273
Heath, J.W., Kidd, G.J., Trapp, B.D. and Dunkley, P.R., Myelin maintenance by Schwann cells in the absence of axons, 277
Henderson, Z., see Sherriff, F.E., 52
Henry, J.L., see Yashpal, K., 269
Herrera-Marschitz, M., see O'Connor, W.T., 66
Heym, C., see Kummer, W., 187
Hirai, Y., see Yamaguchi, T., 90
Holzer, P., see Saria, A., 165
Hönack, D., see Löscher, W., 150
Hönack, D., see Löscher, W., 191
Hooiveld, M.H.W., see Joosten, E.A.J., 25
Hope, B.T., see Pasqualotto, B.A., 155
Hori, T., see Nakashima, T., 97
Hosokawa, K., see Yamaguchi, F., 273
Hosono, R. and Kamiya, Ya., Additional genes which result in an elevation of acetylcholine levels by mutations in *Caenorhabditis elegans*, 243
Hoyer, S., see Nitsch, R., 199
Huber, G. and Schuler, A., Characterization of a new 120 kDa micro-

tubule-associated protein (MAP) of rat brain, 221
Hummel, C., see Thürauf, N., 297
Hunter, L.N., see Gentle, M.J., 113
Hurley, K.M., see Spencer, S.E., 217
Hylden, J.L.K., see Nahin, R.L., 226

Imahori, K., see Ishiguro, K., 195
Inagaki, N., Fukui, H., Ito, S. and Wada, H., Type-2 astrocytes show intracellular Ca^2 elevation in response to various neuroactive substances, 257
Ishiguro, K., Omori, A., Sato, K., Tomizawa, K., Imahori, K. and Uchida, T., A serine/threonine proline kinase activity is included in the tau protein kinase fraction forming a paired helical filament epitope, 195
Ishiura, S., see Kwak, S., 21
Itano, T., see Yamaguchi, F., 273
Ito, M., see Kondo, K., 245
Ito, S., see Inagaki, N., 257
Iwamoto, N. and Emson, P.C., Demonstration of neurofibrillary tangles in parvalbumin-immunoreactive interneurones in the cerebral cortex of Alzheimer-type dementia brain, 81

Janjua, N.A., see Yamaguchi, F., 273
Johnson, D., see Mulligan, L.P., 42
Joh, T.H., see Park, D.H., 240
Joosten, E.A.J., Van der Ven, P.F.M., Hooiveld, M.H.W. and Ten Donkelaar, H.J., Induction of corticospinal target finding by release of a diffusible, chemotropic factor in cervical spinal grey matter, 25
Jouvet, M., see Lin, J.S., 61
Jouvet, M., see Luppi, P.-H., 33

Kamiya, Ya., see Hosono, R., 243
Kano, M., Kano, M.-S. and Maekawa, K., Simple spike modulation of Purkinje cells in the cerebellar nodulus of the pigmented rabbit to optokinetic stimulation, 101
Kano, M.-S., see Kano, M., 101
Kawana, A., see Yamashita, A., 147
Keim, P.S., see Anderson, J.P., 126
Kempf, E., see Lannes, B., 177
Khanna, S.M., see Brundin, Lou, 77
Kibbe, M.R., see Spencer, S.E., 217
Kidd, G.J., see Heath, J.W., 277
Kincaid, A.E., Penny Jr., J.B., Young, A.B. and Newman, S.W., Evidence for a projection from the globus pallidus to the entopeduncular nucleus in the rat, 121
Kitahama, K., see Lin, J.S., 61
Kiyama, Hiroshi, Sato, M., Emson, P.C. and Tohyama, M., Transient expression of neurotensin mRNA in the mitral cells of rat olfactory bulb during development, 85
Kiyohara, T., see Nakashima, T., 97
Kobal, G., see Thürauf, N., 297
Koistinaho, J., Nicotine-induced Fos-like immunoreactivity in rat sympathetic ganglia and adrenal medulla, 47
Kondo, K., Murase, T., Otake, K., Ito, M. and Oiso, Yu., Centrally administered galanin inhibits osmotically stimulated arginine vasopressin release in conscious rats, 245
Koshimura, K., see Ohue, T., 93
Krane, J., see Sonnewald, U., 235
Krause, D., Mischeck, U., Galla, H.J. and Dermietzel, R., Correlation of zonula occludens ZO-1 antigen expression and transendothelial resistance in porcine and rat cultured cerebral endothelial cells, 301
Kummer, W. and Heym, C., Different types of calcitonin gene-related peptide-immunoreactive neurons in the guinea-pig stellate ganglion as revealed by triple-labelling immunofluorescence, 187
Kuraishi, Ya., see Yamaguchi, T., 90
Kwak, S., Masaki, T., Ishiura, S. and Sugita, H., Multicatalytic proteinase is present in Lewy bodies and neurofibrillary tangles in diffuse Lewy body disease brains, 21

Laduron, P.M., see Stutzmann, J.M., 4
Lamour, Y., see Lee, B.H., 29
Lannes, B., Micheletti, G., Warter, J.-M., Kempf, E. and Scala, G.D., Behavioural, pharmacological and biochemical effects of acute and chronic administration of ketamine in the rat, 177
Lee, B.H., Lamour, Y. and Bassant, M.H., Iontophoretic study of medial septal neurons in the unanesthetized rat, 29
Lee, K., see Ohue, T., 93
Lemeignan, M., see Musumeci, D., 285
Levi, G., Agresti, C., D'Urso, D. and Aloisi, F., Is the oligodendroglial differentiation of bipotential oligodendrocyte–type 2 astrocyte progenitors promoted by autocrine factors?, 37
L'Héreault, S. and Barden, N., Biphasic regulation of proopiomelanocortin messenger RNA concentrations by corticotropin-releasing factor in primary cultures of rat hypothalamic neurons, 207
Lieberburg, I., see Anderson, J.P., 126
Lindefors, N., see O'Connor, W.T., 66
Lin, J.S., Fort, P., Kitahama, K., Panula, P., Denney, R.M. and Jouvet, M., Immunohistochemical evidence for the presence of type B monoamine oxidase in histamine-containing neurons in the posterior hypothalamus of cats, 61
Long, S.K., see Wesselman, J.P.M., 261
Löscher, W., Annies, R. and Hönack, D., The *N*-methyl-D-aspartate receptor antagonist MK-801 induces increases in dopamine and serotonin metabolism in several brain regions of rats, 191
Löscher, W., Hönack, D. and Taylor, C.P., Gabapentin increases aminooxyacetic acid-induced GABA accumulation in several regions of rat brain, 150
Louw, D., see Sutherland, G., 169
Lucas, M., see Stutzmann, J.M., 4
Lundeberg, T., Eriksson, S.V. and Theodorsson, E., Neuroimmunomodulatory effects of acupuncture in mice, 161
Luppi, P.-H., Charlety, P.J., Fort, P., Akaoka, H., Chouvet, G. and Jouvet, M., Anatomical and electrophysiological evidence for a glycinergic inhibitory innervation of the rat locus coeruleus, 33

Maekawa, K., see Kano, M., 101
Maggi, C.A., see Saria, A., 165
Marlier, L., Poulat, P., Rajaofetra, N. and Privat, A., Modification of serotonergic immunoreactive pattern in the dorsal horn of the rat spinal cord following dorsal root rhizotomy, 9
Maršala, J., see Orendáčová, J., 143
Maršala, M., see Orendáčová, J., 143
Masaki, T., see Kwak, S., 21
Mazurkiewicz, J.E., Ubiquitin deposits are present in spinal motor neurons in all stages of the disease in the motor neuron degeneration (*Mnd*) mutant of the mouse, 182
McBride, R.L., see Garrett, W.T., 137
McGregor, G.P., see Anand, P., 57
Micheletti, G., see Lannes, B., 177
Mileson, B.E. and Schwartz, R.D., The use of locomotor activity as a behavioral screen for neuronal damage following transient forebrain ischemia in gerbils, 71
Minami, M., see Yamaguchi, T., 90
Mischeck, U., see Krause, D., 301
Misgeld, U., see Beyer, C., 1
Miwa, S., see Ohue, T., 93

Miyamoto, O., see Yamaguchi, F., 273
Mizuno, N., see Al-Yousuf, S., 249
Morrison, J.F.B., see Anand, P., 57
Morrison, J.F.B., see Sherriff, F.E., 52
Mulligan, L.P., Rosenblatt, D.E., Toms, R. and Johnson, D., Protease nexin-1 activity in cultured Schwann cells, 42
Murase, T., see Kondo, K., 245
Musumeci, D., Lemeignan, M. and Bloch, S., Interocular transfer of depth discrimination in intact and in DSO-sectioned pigeons, 285

Nahin, R.L. and Hylden, J.L.K., Peripheral inflammation is associated with increased glutamic acid decarboxylase immunoreactivity in the rat spinal cord, 226
Naik, B., see Silverstein, F.S., 133
Nakai, S., see Yamaguchi, T., 90
Nakashima, T., Kiyohara, T. and Hori, T., Tumor necrosis factor-β specifically inhibits the activity of preoptic warm-sensitive neurons in tissue slices, 97
Needleman, P., see Spencer, S.E., 217
Newman, S.W., see Kincaid, A.E., 121
Nitsch, R. and Hoyer, S., Local action of the diabetogenic drug, streptozotocin, on glucose and energy metabolism in rat brain cortex, 199

O'Connor, W.T., Lindefors, N., Brené, S., Herrera-Marschitz, M., Persson, H. and Ungerstedt, Urban, Short-term dopaminergic regulation of GABA release in dopamine deafferented caudate-putamen is not directly associated with glutamic acid decarboxylase gene expression, 66
O'Donovan, M.J., see Sernagor, E., 211
Ögren, S.O. and Pramanik, A., Galanin stimulates acetylcholine release in the rat striatum, 253
Ohashi, Ya., see Ando, S., 17
Ohmoto, T., see Yamaguchi, F., 273
Ohue, T., Koshimura, K., Lee, K., Watanabe, Ya. and Miwa, S., A novel action of 6*R*-L-*erythro*-5,6,7,8-tetrahydrobiopterin, a cofactor for hydroxylases of phenylalanine, tyrosine and tryptophan: enhancement of acetylcholine release in vivo in the rat hippocampus, 93
Oiso, Yu., see Kondo, K., 245
Omori, A., see Ishiguro, K., 195
Orendáčová, J., Maršala, M. and Maršala, J., The blood–brain barrier permeability in graded postischemic spinal cord reoxygenation in rabbits, 143
Otake, K., see Kondo, K., 245

Panula, P., see Lin, J.S., 61
Paran, Y., see Seltzer, Z., 203
Park, D.H., Park, H.S., Joh, T.H., Anwar, M. and Ruggiero, D.A., Strain difference in phenylethanolamine *N*-methyltransferase activity and immunoreactivity of medulla oblongata of Sprague–Dawley and Long–Evans hooded rats, 240
Park, H.S., see Park, D.H., 240
Pasqualotto, B.A., Hope, B.T. and Vincent, S.R., Citrulline in the rat brain: immunohistochemistry and coexistence with NADPH-diaphorase, 155
Penny Jr., J.B., see Kincaid, A.E., 121
Persson, H., see O'Connor, W.T., 66
Petersen, S.B., see Sonnewald, U., 235
Petrig, B.L., see Riva, C.E., 291
Pilgrim, C., see Beyer, C., 1
Pinsky, C., see Sutherland, G., 169
Piot, O., see Stutzmann, J.M., 4
Poulat, P., see Marlier, L., 9
Pramanik, A., see Ögren, S.O., 253
Privat, A., see Marlier, L., 9

Radhakrishnan, V., see Yashpal, K., 269
Rajaofetra, N., see Marlier, L., 9
Randall, W.C., see Xi, X., 129
Rataud, J., see Stutzmann, J.M., 4
Reisert, I., see Beyer, C., 1
Reuss, S., Photoperiod effects on bombesin- and cholecystokinin-like immunoreactivity in the suprachiasmatic nuclei of the Djungarian hamster (*Phodopus sungorus*), 13
Richmond, S., see Brammer, M.J., 231
Riva, C.E., Harino, S., Shonat, R.D. and Petrig, B.L., Flicker evoked increase in optic nerve head blood flow in anesthetized cats, 291
Robakis, N.K., see Anderson, J.P., 126
Rosenblatt, D.E., see Mulligan, L.P., 42
Ruggiero, D.A., see Park, D.H., 240

Sambamurti, K., see Anderson, J.P., 126
Saper, C.B., see Spencer, S.E., 217
Saria, A., Maggi, C.A. and Holzer, P., Functional and pharmacological aspects of neuropeptides: a conference report of the first meeting of the European Neuropeptide Club, 165
Satoh, Masamichi, see Yamaguchi, T., 90
Sato, K., see Ishiguro, K., 195
Sato, M., see Kiyama, Hiroshi, 85
Scala, G.D., see Lannes, B., 177
Scaravilli, F., see Anand, P., 57
Schousboe, A., see Sonnewald, U., 235
Schuler, A., see Huber, G., 221
Schwartz, R.D., see Mileson, B.E., 71
Seiger, Å., see Hao, J. -X., 105
Seltzer, Z., Paran, Y., Eisen, A. and Ginzburg, R., Neuropathic pain behavior in rats depends on the afferent input from nerve-end neuroma including histamine-sensitive C-fibers, 203
Sernagor, E. and O'Donovan, M.J., Whole-cell patch clamp recordings from rhythmically active motoneurons in the isolated spinal cord of the chick embryo, 211
Sherriff, F.E., Henderson, Z. and Morrison, J.F.B., Further evidence for the absence of a descending cholinergic projection from the brainstem to the spinal cord in the rat, 52
Shonat, R.D., see Riva, C.E., 291
Silva, M.A., see Brammer, M.J., 231
Simoncsits, A., see Bristulf, J., 173
Silverstein, F.S. and Naik, B., Effect of depolarization on striatal amino acid efflux in perinatal rats: an in vivo microdialysis study, 133
Sonnewald, U., Westergaard, N., Krane, J., Unsgård, G., Petersen, S.B. and Schousboe, A., First direct demonstration of preferential release of citrate from astrocytes using [^{13}C]NMR spectroscopy of cultured neurons and astrocytes, 235
Spencer, S.E., Kibbe, M.R., Hurley, K.M., Needleman, P. and Saper, C.B., Origin of porcine brain natriuretic peptide-like immunoreactive innervation of the middle cerebral artery in the rat, 217
Strijkstra, A.M., see Daan, S., 265
Stutzmann, J.M., Eon, B., Darche, F., Lucas, M., Rataud, J., Piot, O., Blanchard, J.C. and Laduron, P.M., Are 5-HT$_2$ antagonists endowed with anxiolytic properties in rodents?, 4
Sugita, H., see Kwak, S., 21
Sutherland, G., Bose, R., Louw, D. and Pinsky, C., Global elevation of brain superoxide dismutase activity following forebrain ischemia in rat, 169
Sveinsson, K., see Altmann, L., 109

Tagliavini, F., Giaccone, G., Verga, L., Ghiso, J., Frangione, B. and Bugiani, O., Alzheimer patients: preamyloid deposits are immunoreactive with antibodies to extracellular domains of the amyloid precursor protein, 117
Taylor, C.P., see Löscher, W., 150
Ten Donkelaar, H.J., see Joosten, E.A.J., 25
Theodorsson, E., see Lundeberg, T., 161
Thürauf, N., Friedel, I., Hummel, C. and Kobal, G., The mucosal potential elicited by noxious chemical stimuli with CO_2 in rats: is it a peripheral nociceptive event?, 297
Tohyama, M., see Kiyama, Hiroshi, 85
Tomizawa, K., see Ishiguro, K., 195
Toms, R., see Mulligan, L.P., 42
Trapp, B.D., see Heath, J.W., 277

Uchida, T., see Ishiguro, K., 195
Ulfendahl, M., see Brundin, Lou, 77
Ungerstedt, Urban, see O'Connor, W.T., 66
Unsgård, G., see Sonnewald, U., 235

Van der Ven, P.F.M., see Joosten, E.A.J., 25
Van Wilgenburg, H., see Wesselman, J.P.M., 261
Verga, L., see Tagliavini, F., 117
Vincent, S.R., see Pasqualotto, B.A., 155

Wada, H., see Inagaki, N., 257
Waddington, D., see Gentle, M.J., 113
Warter, J.-M., see Lannes, B., 177
Watanabe, Ya., see Ohue, T., 93
Wesselman, J.P.M., Van Wilgenburg, H. and Long, S.K., The effects of pentobarbital and benzodiazepines on GABA-responses in the periphery and spinal cord in vitro, 261
Westergaard, N., see Sonnewald, U., 235
Wiegand, H., see Altmann, L., 109
Wiesenfeld-Hallin, Z., see Hao, J.-X., 105
Williams Jr., J.K., see Garrett, W.T., 137
Wright, A.K., Arbuthnott, G.W. and Dunnett, S.B., Serotonin hyperinnervation after foetal nigra or raphe transplantation in the neostriatum of adult rats, 281
Wurster, R.D., see Xi, X., 129

Xiang, J.Z., see Brammer, M.J., 231
Xi, X., Randall, W.C. and Wurster, R.D., Intracellular recording of spontaneous activity of canine intracardiac ganglion cells, 129
Xu, X.-J., see Hao, J.-X., 105

Yamaguchi, F., Itano, T., Miyamoto, O., Janjua, N.A., Ohmoto, T., Hosokawa, K. and Hatase, O., Increase of extracellular insulin-like growth factor I (IGF-I) concentration following electrolytical lesion in rat hippocampus, 273
Yamaguchi, T., Kuraishi, Ya., Minami, M., Nakai, S., Hirai, Y. and Satoh, Masamichi, Methamphetamine-induced expression of interleukin-1β mRNA in the rat hypothalamus, 90
Yamashita, A. and Kawana, A., Nerve growth factor-induced intracellular calcium ion release in chick dorsal root ganglion neurons, 147
Yashpal, K., Radhakrishnan, V. and Henry, J.L., NMDA receptor antagonist blocks the facilitation of the tail flick reflex in the rat induced by intrathecal administration of substance P and by noxious cutaneous stimulation, 269
Young, A.B., see Kincaid, A.E., 121
Yu, Y.-X., see Hao, J.-X., 105